REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
Public reporting burden for this maintaining the data needed a	collection of informatio	n is estimated to average 1 hour	per response, including the time for	reviewing instructions, s	searching existing data sources, gathering and	
including suggestions for reduc	ring this hurden to Done	ertment of Defense, Westington	in. Send comments regarding this t	ourden estimate or any o	ther aspect of this collection of information.	
righway, Suite 1204, Arlingtor collection of information if it do	ı, VA 22202-4302. Res es not display a currentl	pondents should be aware that r	notwithstanding any other provision EASE DO NOT RETURN YOUR F	of law, no person shall b	ns and Reports (0704-0188), 1215 Jefferson Davi e subject to any penalty for failing to comply with	
1. REPORT DATE (DE	-MM-YYYY)	2. REPORT TYPE	LASE DO NOT RETURN TOUR FO		3. DATES COVERED (From - To)	
	•	Technical Papers	;	1	3. DATES COVERED (From - 10)	
4. TITLE AND SUBTIT	LE				5a. CONTRACT NUMBER	
				[:	
				ļ-	5b. GRANT NUMBER	
					!	
					c. PROGRAM ELEMENT NUMBER	
					· · · · · · · · · · · · · · · · · · ·	
6. AUTHOR(S)					od. PROJECT NUMBER	
				(2303 :	
				<u></u>	ランクラ 5e. TASK NUMBER	
			,	<u></u>	mac8	
		•		5	f. WORK UNIT NUMBER	
7 PERFORMING ORG	ANIZATION NAM	E(S) AND ADDRESS(E				
7. I EM ONWING ONG	ANIZATION NAM	E(S) AND ADDRESS(E	S) ·		3. PERFORMING ORGANIZATION	
Air Force Research I	aboratory (AEN	AC)		F	REPORT	
AFRL/PRS	abbratory (AFF	vic)				
5 Pollux Drive						
Edwards AFB CA 9	2524 7040					
Edwards AFB CA 9.	3524-7048				•	
					4	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)					0. SPONSOR/MONITOR'S	
			` ,		CRONYM(S)	
= -						
Air Force Research L	aboratory (AFN	(IC)				
AFRL/PRS				1	1. SPONSOR/MONITOR'S	
5 Pollux Drive				1 -	NUMBER(S)	
Edwards AFB CA 93	524-7048					
12. DISTRIBUTION / AV	AU ADU ITV OTA					
12. DISTRIBUTION / AV	ALABILITY STA	IEMENI				
Approved for public r	برادينا المعالم	* 31 1. 1			•	
Approved for public r	elease; distribut	ion unlimited.				
13. SUPPLEMENTARY	NOTES					
					•	
14. ABSTRACT						
					0	
15 CUD IEOT TERMS						
15. SUBJECT TERMS				2.0		
			1 -	22 0	20	
IC CECUDITY OF ACCU	OATION				-	
16. SECURITY CLASSIF	CATION OF:		17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE	
			OF ABSTRACT	OF PAGES	PERSON	
DEDORT	ADOTO				Leilani Richardson	
a. REPORT b	ABSTRACT	c. THIS PAGE	,		19b. TELEPHONE NUMBER	
Including T	Inclosed 1	¥7 3 400 -	(A)	1	(include area code)	
Unclassified U	nclassified	Unclassified	1 \		(661) 275-5015	

Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. 239.18

MEMORANDUM FOR PRS

FROM: PROI (TI) (STINFO)

28 May 1999

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-FY99-0117 Yoo and Boatz, "Theoretical Study of Decomposition Mechanism of High Energy Density Materials"

Poster Session HEDM Conference

(Statement A)

020 20021122

Hi Young Yoo and Jerry Boatz*

Theoretical Study of Decomposition Mechanism of

High Energy Density Materials

*Air Force Research Laboratory, Edwards AFB, CA 93524-7680 National Research Council, Washington, DC 20418

AFOSR HEDM CONTRACTOR'S MEETING Cocoa Beach, FL 6/8-6/10, 1999

DISTRIBUTION STATEMENT A: Approved for Public Release -**Distribution Unlimited**

Abstract

Decomposition mechanisms of [NH₂Me₂]⁺[NO₃], and [NH₂(NH₂)₂]⁺[NO₃] were processes have been explored: (1) proton transfer and (2) Me⁺/NH₂⁺ transfer 31G* level and single point energies at the MP2/6-31G*//RHF/6-31G* level have been calculated. Comparison of similarities and differences between these two systems will be presented here. Future research will involve characterization of the potential energy surface for [NH2NMe2NH2]*[NO3] and design of potential catalysts One of the HEDM research efforts has focused on the development of high performance monopropellants in replacement for hydrazine due to its toxicity. A stabilities of monopropellants by using ab initio quantum mechanical calculations. investigated. The potential energy surfaces of two gas-phase decomposition reactions. Transition states for both pathways have been located at the RHF/6proposed area of research involves determination of decomposition mechanisms and that will stabilize transition states.

$$[NH_2Me_2]^+[NO_3]^- \rightarrow NHMe_2 + HNO_3$$
 (1a)

$$[NH_2Me_2]^+[NO_3]^- \rightarrow NH_2Me + MeNO_3$$
 (2a)

$$[NH_2(NH_2)_2]^+[NO_3]^- \to NH(NH_2)_2 + HNO_3$$
 (1b)

$$[NH_2(NH_2)_2]^+[NO_3]^- \to NH_2NH_2 + NH_2NO_3$$
 (2b)

Background

Hydrazine is the state of the art monopropellant currently. However, it has several disadvantages including toxicity, volatility, and handling. Researchers are continually looking for new monopropellant candidates which include energetic materials such as substituted ammonium salts. Our research effort has concentrated on the decomposition routes of some HEDM materials. Ab initio quantum mechanical calculations have been carried out by our group to explore possible decomposition mechanisms.

The model compounds under study are $[NH_2Me_2]^+[X]^-$, where $X = NO_3$ and CI, and $[N(NH_2)_2Me_2]^+[NO3]$.

Objectives:

- To map out the potential energy surface of these model compounds
 - To determine the energetics of decomposing species
- To design a catalyst that stabilizes the transition state

Gaseous products detected from experimental thermal decomposition studies of $N(NH_2)_2(Me_2)NO_3$ are N_2 , NH_3 , H_2O and CH_4 .

Possible reaction pathways leading to the final products are the following: R, R' = H, CH₃; X= NO₃

(1) $[N(NH_2)_2RR']^+[X]^- \rightarrow [N(NH_2)_2R'][RX]$ (1b) $[N(NH_2)_2RR']^+[X]^- \rightarrow [N(NH_2)RR'][NH_2X]$

(2a) $N(NH_2)_2R' \rightarrow N(NH_2)R' + NH_2$ (2b) $N(NH_2)RR' \rightarrow NRR' + NH_2$ (3) $[N(NH_2)_2RR']^+ \rightarrow RH + [NH_3R']^+ + N_2$

(4a) $NH_2X \rightarrow NH_2 + X$

4b) $NH_2 + NH_2X \rightarrow NH_3 + NHX$

(4c) $X + NH_2X \rightarrow HX + NHX$

(4d) 2NHX \rightarrow X-NH-NH-X

(4e) X-NH-NH-X \rightarrow 2HX + N₂

Rubtsov, Y. I.; Andrienko, L. P.; Titova, K. V.; Loginova, E. N. Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya 1982, 9, 1953.

Decomposition Mechanisms of [NH₂(NH₂)₂]⁺[X]⁻, $X = CI \text{ or } NO_3$


Proton Transfer


$$[NH_2(NH_2)_2]^+[X]^- \rightarrow NH(NH_2)_2 + HX$$

NH2 Cation Transfer

$$[NH_2(NH_2)_2]^+[X]^- \rightarrow NH_2NH_2 + NH_2X$$

A Proton Transfer Transition Structure of [NH₂(NH₂)₂]⁺[NO₃] At RHF/6-31G*(6d)






Intermediate Complex


Energy 0.0 (kcal/mol) (0.0)

Transition Structure

1.3

A NH2⁺ Transfer Transition Structure of [NH2(NH2)2]⁺[NO3] At RHF/6-31G*(6d)





Intermediate Complex


Energy (kcal/mol)

0.0 (0.0)


Transition Structure


59.8

(55.9)



A Proton Transfer Transition Structure of [NH₂Me₂]⁺[NO₃] At RHF/6-31G*(6d)





Intermediate Complex

0.0

(kcal/mol)


Energy


Transition Structure

2.2

(-1.3)

A Me⁺ Transfer Transition Structure of [NH₂Me₂]⁺[NO₃]⁻ At RHF/6-31G*(6d)






Intermediate Complex

Energy (kcal/mol)

 $0.0 \\ (0.0)$

Transition Structure

45.8 (47.7)



Conclusion

Based on the ab initio calculation studies of [NH₂(NH₂)₂]⁺[NO₃ or Cl] and $[\mathrm{NH_2(Me)_2}]^+[\mathrm{NO_3} \text{ or CI}]^-$

- The transition structure of proton transfer is earlier than that of methyl(or NH_2) cation transfer and also lower in barrier.
- Variations on anions or cations did not affect on the energy barrier or the transition structure.
- calculations imply that methyl cation transfer of [NMe₂(NH₂)₂]⁺[NO₃]⁻ will Since variations on cations did not affect on the energy barrier, our be favorable than NH₂⁺.

Future Work

- The decomposition mechanism of $[NMe_2(NH_2)_2]^+[NO_3]^-$ as well as other ammonium salt monopropellant candidates will be investigated theoretically.
- potential energy surface of [NMe₂(NH₂)₂]⁺[NO₃] and it will be compared Different catalysts will be included in the calculations for determining the with the uncatalyzed potential energy surface.